

## EURO-PERISTAT DATA COLLECTION INSTRUMENT

## CORE INDICATORS OF PERINATAL HEALTH

This data collection instruments collects data on perinatal health indicators for the EURO-PERISTAT project. These data will be used to develop recommendations for a European perinatal health monitoring system by the EURO-PERISTAT Scientific Committee.

These data will not be made public in reports or publications without advanced approval from respondents. Respondents and their institutions will be formally acknowledged in all documents.

## General Instructions

1. To run this Excel file correctly, the Macro security of the Excel program should be set to 'medium' (Go to Tools -> Macro -> Security). This should be done before starting the Peristatcore.xls file.
2. Please provide data for the year **2010**, or most recent year available **before** 2010.
3. We would like **COUNTRY**-level data. If country-level data are not available, but **POPULATION**-based data from one or more regions are, please use this source.
4. If several data sources are used (e.g. merged data from two sources), please create a new data source sheet and describe each data source in this sheet.
5. If data are provided for different regions within one country please copy and fill in one Peristatcore.xls file **per region**. However, always include country-level data if available.
6. For each table, please register the name of the data source you have used. For each source of data, please complete a data source information form (push button 'to data source information form' on the INDEX sheet OR button 'new data source' on each indicator sheet).
7. For items not defined explicitly in the text use the WHO recommended definitions (push button): WHO ICD-10 Definitions
8. We are specifically requesting that you provide us **WITH NUMBERS**. Rates will be calculated based on the numbers you provide.
9. All tables for **Core** Indicators can be accessed from the INDEX sheet by clicking on the corresponding 'Go to indicator' button.
10. In the tables totals will be calculated automatically. Please use these as a check.
11. After filling in a table and returning to the index sheet via the button 'Save and return to INDEX' the box of the corresponding table will be ticked, indicating that data on this indicator have been filled in.
12. For more detailed instructions on this Data Collection Instrument please see the Instruction Manual.

[To INDEX sheet](#)

Index sheet **CORE** Indicators

To data source information form

<b>Core Indicator # 0</b>			Data source used:
Recording of births and deaths	<input type="checkbox"/>	Go to indicator	<Unknown>
<b>Core Indicator # 1</b>			
Fetal mortality by gestational age	<input type="checkbox"/>	Go to indicator	<Unknown>
Fetal mortality by birth weight	<input type="checkbox"/>	Go to indicator	<Unknown>
<b>Core Indicator # 2</b>			
Neonatal mortality by gestational age (annual)	<input type="checkbox"/>	Go to indicator	<Unknown>
Neonatal mortality by birth weight (annual)	<input type="checkbox"/>	Go to indicator	<Unknown>
Neonatal mortality by gestational age (cohort)	<input type="checkbox"/>	Go to indicator	<Unknown>
Neonatal mortality by birth weight (cohort)	<input type="checkbox"/>	Go to indicator	<Unknown>
<b>Core Indicator # 3</b>			
Infant mortality by gestational age (annual)	<input type="checkbox"/>	Go to indicator	<Unknown>
Infant mortality by birth weight (annual)	<input type="checkbox"/>	Go to indicator	<Unknown>
Infant mortality by gestational age (cohort)	<input type="checkbox"/>	Go to indicator	<Unknown>
Infant mortality by birth weight (cohort)	<input type="checkbox"/>	Go to indicator	<Unknown>
<b>Core Indicator # 4</b>			
Distribution of birth weight by plurality	<input type="checkbox"/>	Go to indicator	<Unknown>
Distribution of birth weight by gestational age	<input type="checkbox"/>	Go to indicator	<Unknown>
<b>Core Indicator # 5</b>			
Distribution of gestational age by plurality	<input type="checkbox"/>	Go to indicator	<Unknown>
<b>Core Indicator # 6</b>			
Maternal mortality by maternal age (routine)	<input type="checkbox"/>	Go to indicator	<Unknown>
Maternal mortality by maternal age (enhanced)	<input type="checkbox"/>	Go to indicator	<Unknown>
<b>Core Indicator # 7</b>			
Multiple maternity rates	<input type="checkbox"/>	Go to indicator	<Unknown>
<b>Core Indicator # 8</b>			
Distribution of maternal age	<input type="checkbox"/>	Go to indicator	<Unknown>
<b>Core Indicator # 9</b>			
Distribution of parity	<input type="checkbox"/>	Go to indicator	<Unknown>
<b>Core Indicator # 10</b>			
Mode of delivery	<input type="checkbox"/>	Go to indicator	<Unknown>
Mode of delivery by parity	<input type="checkbox"/>	Go to indicator	<Unknown>
Mode of delivery by previous Caesarean	<input type="checkbox"/>	Go to indicator	<Unknown>
Mode of delivery by presentation	<input type="checkbox"/>	Go to indicator	<Unknown>
Mode of delivery by plurality	<input type="checkbox"/>	Go to indicator	<Unknown>
Mode of delivery by gestational age	<input type="checkbox"/>	Go to indicator	<Unknown>

To data source information form

**Core indicator #6: Maternal mortality (by maternal age)**

**Definition:** The number of maternal deaths (the death of a woman while pregnant or within 42 days of the termination of pregnancy, irrespective of the duration and site of the pregnancy, for any cause related to or aggravated by the pregnancy or its management, but not from accidental or incidental causes) per 100 000 live births using data from routine cause of death statistics.

SAVE and  
return to INDEX

WHO definition maternal death

Can you provide data using another definition? ☐ yes ☐ no

Please identify any routine enhanced registration or casefinding methods for maternal mortality in your country (confidential enquiry, routine linkage of vital records):

☐ yes ☐ no

### New data source

	Year 1	Year 2	Year 3	Year 4	Year 5
Give data for 5 consecutive years (pref. 2006-2007-2008-2009-2010):					

[Reset this page](#)

None

Maternal age	Year 1		2		Year 3		4		5	
	N maternal deaths	N live births	Year	N maternal	N live births	N maternal deaths	N live births	Year	N maternal	N live births
10-14										
15-19										
20-24										
25-29										
30-34										
35-39										
40-44										
45-49										
50+										
Unknown										
Calculated totals	0	0		0	0		0		0	0



### Core indicator #6: Maternal mortality (by maternal age)

**Definition:** The number of maternal deaths (the death of a woman while pregnant or within 42 days of the termination of pregnancy, irrespective of the duration and site of the pregnancy, for any cause related to or aggravated by the pregnancy or its management, but not from accidental or incidental causes) per 100 000 live births using data from routine cause of death statistics.

SAVE and  
return to INDEX

*Attention: only include women delivering live and stillbirths at or after 22 completed weeks of gestation.*

WHO definition maternal death

Are you able to provide data using this definition? ☒ yes ☐ no  
Can you provide data using another definition? ☒ yes ☐ no

Is there an indicator of pregnancy (e.g. a checkbox) on the death certificate in your country? ☒ yes ☐ no

Please identify any routine enhanced registration or casefinding methods for maternal mortality in your country (confidential enquiry, routine linkage of vital records):

To INDEX  
without saving

In the table below all maternal deaths within 42 days after termination of pregnancy are included. For future purposes, could you also provide maternal deaths up to 1 year after termination of pregnancy? ☒ yes ☐ no

Data source:

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Give data for 5 consecutive years (pref. 2006-2007-2008-2009-2010):

Year 1	Year 2	Year 3	Year 4	Year 5
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

Comments:

Reset this page

Select one of  
these options:

☒ option 1 (preferred)

☐ option 2 (totals only)

☐ None

Maternal age

Totals

Year 1		Year 2		Year 3		Year 4		Year 5	
N maternal deaths	N live births	N maternal deaths	N live births	N maternal deaths	N live births	N maternal deaths	N live births	N maternal deaths	N live births
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
Totals									

### Core indicator #6: Maternal mortality (by maternal age)

**Definition:** The number of maternal deaths (the death of a woman while pregnant or within 42 days of the termination of pregnancy, irrespective of the duration and site of the pregnancy, for any cause related to or aggravated by the pregnancy or its management, but not from accidental or incidental causes) per 100 000 live births using data from routine cause of death statistics.

SAVE and  
return to INDEX

WHO definition maternal death

*Attention: only include women delivering live and stillbirths at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☒ yes ☐ no  
Can you provide data using another definition? ☐ yes ☐ no

Is there an indicator of pregnancy (e.g. a checkbox) on the death certificate in your country? ☒ yes ☐ no

Please identify any routine enhanced registration or casefinding methods for maternal mortality in your country (confidential enquiry, routine linkage of vital records):

In the table below all maternal deaths within 42 days after termination of pregnancy are included. For future purposes, could you also provide maternal deaths up to 1 year after termination of pregnancy? ☒ yes ☐ no

Data source: 0 New data source

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Give data for 5 consecutive years (pref. 2006-2007-2008-2009-2010):

Year 1	Year 2	Year 3	Year 4	Year 5

Comments:

To INDEX  
without saving

Reset this page

Select one of  
these options:

☒ option 1 (preferred)

☐ option 2 (totals only)

☐ None

If you cannot provide data on this indicator using the categories used in the proposed options, please send us a separate Excel table with data as reported in your country.  
In this Excel table please clearly specify the indicator, used categories and row and column headers.

### Core indicator #6: Maternal mortality (by maternal age)

**Definition:** The number of maternal deaths (the death of a woman while pregnant or within 42 days of the termination of pregnancy, irrespective of the duration and site of the pregnancy, for any cause related to or aggravated by the pregnancy or its management, but not from accidental or incidental causes) per 100 000 live births using data from enhanced systems.

WHO definition maternal death

**Attention:** only include women delivering live and stillbirths at or after 22 completed weeks of gestation.

Are you able to provide data using this definition? ☐ yes ☐ no

Can you provide data using another definition? ☐ yes ☐ no

Is there a report published from this enhanced system? ☐ yes ☐ no

In the table below all maternal deaths within 42 days after termination of pregnancy are included. For future purposes, could you also provide maternal deaths up to 1 year after termination of pregnancy? ☐ yes ☐ no

Data source:

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Give data for 5 consecutive years (pref. 2006-2007-2008-2009-2010):

Year 1	Year 2	Year 3	Year 4	Year 5
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

Comments:

Select one of these options:

☒ option 1 (preferred)

☐ option 2 (totals only)

☐ None

Maternal age	Year 1		Year 2		Year 3		Year 4		Year 5	
	N maternal deaths	N live births	N maternal deaths	N live births	N maternal deaths	N live births	N maternal deaths	N live births	N maternal deaths	N live births
10-14										
15-19										
20-24										
25-29										
30-34										
35-39										
40-44										
45-49										
50+										
Unknown										
Calculated totals	0	0	0	0	0	0	0	0	0	0

[SAVE and return to INDEX](#)

[To INDEX without saving](#)

[Reset this page](#)

SAVE and  
return to INDEX

WHO definition maternal death

Are you able to provide data using this definition? ☐ yes ☐ no

Can you provide data using another definition? ☐ yes ☐ no

Is there a report published from this enhanced system? ☒ yes ☐ no

To INDEX  
without saving

In the table below all maternal deaths within 42 days after termination of pregnancy are included. For future purposes, could you also provide maternal deaths up to 1 year after termination of pregnancy?

Data source: 0 New data source

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

	Year 1	Year 2	Year 3	Year 4	Year 5
Give data for 5 consecutive years (pref. 2006-2007-2008-2009-2010):					

Comments:

[Reset this page](#)

Select one of these options:

- option 1 (preferred)

**option 2 (totals only)**

☐ None

Year 1		2		Year 3		4		5	
N maternal deaths	N live births	Year	N maternal	N live births	N maternal deaths	N live births	Year	N maternal	N live births

### Maternal age

Totals



### Core indicator #6: Maternal mortality (by maternal age)

**Definition:** The number of maternal deaths (the death of a woman while pregnant or within 42 days of the termination of pregnancy, irrespective of the duration and site of the pregnancy, for any cause related to or aggravated by the pregnancy or its management, but not from accidental or incidental causes) per 100 000 live births using data from enhanced systems.

WHO definition maternal death

*Attention: only include women delivering live and stillbirths at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☒ yes ☐ no  
Can you provide data using another definition? ☒ yes ☐ no

Is there a report published from this enhanced system? ☐ yes ☒ no

In the table below all maternal deaths within 42 days after termination of pregnancy are included. For future purposes, could you also provide maternal deaths up to 1 year after termination of pregnancy? ☒ yes ☐ no

Data source:

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Give data for 5 consecutive years (pref. 2006-2007-2008-2009-2010):  
Year 1  Year 2  Year 3  Year 4  Year 5

Comments:

Select one of these options:

☒ option 1 (preferred)

☐ option 2 (totals only)

☐ None

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

If you cannot provide data on this indicator using the categories used in the proposed options, please send us a separate Excel table with data as reported in your country.  
In this Excel table please clearly specify the indicator, used categories and row and column headers.



### Core indicator #7: Multiple maternity rate (by number of fetuses)

**Definition:** The number of women in a multiple gestation pregnancy at delivery as a proportion of all women delivering live or stillborn babies.

**Attention:** only include women delivering live and stillbirths at or after 22 completed weeks of gestation.

Are you able to provide data using this definition?

☒ yes

☐ no

Can you provide data using another definition?

☒ yes

☐ no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good

☐ some concerns

☐ bad

How are multiple pregnancies with different outcomes (live + stillbirth) registered?:

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these options:

☒ option 1 (preferred)

☐ option 2

☐ None

N women delivering  
live or stillborn babies

Number of fetuses at delivery

Singleton

Twin

Triplet

Quadruplet+

Unknown

Calculated total

0

### Core indicator #7: Multiple maternity rate (by number of fetuses)

**Definition:** The number of women in a multiple gestation pregnancy at delivery as a proportion of all women delivering live or stillborn babies.

**Attention:** only include women delivering live and stillbirths at or after 22 completed weeks of gestation.

Are you able to provide data using this definition? ☒ yes ☐ no

Can you provide data using another definition? ☐ yes ☒ no

Data source:

[New data source](#)

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

How are multiple pregnancies with different outcomes (live + stillbirth) registered?

Comments:

Select one of these options:

☒ option 1 (preferred)

☐ option 2

☐ None

N women delivering  
live or stillborn babies

#### Number of fetuses at delivery

Singleton

Twin

Triplet+

Unknown

Calculated total

0

[SAVE and  
return to INDEX](#)

[To INDEX  
without saving](#)

[Reset this page](#)

Core indicator #7: Multiple maternity rate (by number of fetuses)		SAVE and return to INDEX
<b>Definition:</b> The number of women in a multiple gestation pregnancy at delivery as a proportion of all women delivering live or stillborn babies. <i>Attention: only include <u>women</u> delivering live and stillbirths at or after 22 completed weeks of gestation.</i>		
Are you able to provide data using this definition? <input type="checkbox"/> yes <input type="checkbox"/> no Can you provide data using another definition? <input type="checkbox"/> yes <input type="checkbox"/> no		
<div style="border: 1px solid black; height: 40px; width: 100%;"></div>		To INDEX without saving
Data source: <span style="border: 1px solid black; padding: 2px;">0</span> <span style="float: right;">New data source</span>		
Please rate the quality of this indicator: <input checked="" type="checkbox"/> good <input type="checkbox"/> some concerns <input type="checkbox"/> bad		Reset this page
How are multiple pregnancies with different outcomes (live + stillbirth) registered?: <div style="border: 1px solid black; height: 40px; width: 100%;"></div>		
Comments: <div style="border: 1px solid black; height: 40px; width: 100%;"></div>		
Select one of these options: <input checked="" type="radio"/> option 1 (preferred) <input type="radio"/> option 2 <input type="radio"/> None		

If you cannot provide data on this indicator using the categories used in the proposed options, please send us a separate Excel table with data as reported in your country. In this Excel table please clearly specify the indicator, used categories and row and column headers.



Core indicator #8: Distribution of maternal age	
<b>Definition:</b> Age in years at delivery for women delivering live or stillborn babies.	
Are you able to provide data using this definition?	<input type="checkbox"/> yes <input type="checkbox"/> no
Can you provide data using another definition?	<input type="checkbox"/> yes <input type="checkbox"/> no
<div></div>	
Data source: 0	New datasource
Please rate the quality of this indicator: <input checked="" type="checkbox"/> good <input type="checkbox"/> some concerns <input type="checkbox"/> bad	
Comments:	

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of  
these options:

☒ option 1 (preferred)

☐ None

N women delivering  
live or stillborn babies

Maternal age

<15	
15-17	
18-19	
20-24	
25-29	
30-34	
35-39	
40-44	
45-49	
50+	
Unknown	

Calculated total 0

Core indicator #8: Distribution of maternal age		SAVE and return to INDEX
<b>Definition:</b> Age in years at delivery for women delivering live or stillborn babies.		
Are you able to provide data using this definition? <input checked="" type="checkbox"/> yes <input type="checkbox"/> no Can you provide data using another definition? <input checked="" type="checkbox"/> yes <input type="checkbox"/> no		To INDEX without saving
<div> <div></div> <div></div> </div>		
Data source: 0 <span>New datasource</span>		Reset this page
Please rate the quality of this indicator: <input checked="" type="checkbox"/> good <input type="checkbox"/> some concerns <input type="checkbox"/> bad		
Comments: <div></div>		
Select one of these options: <input checked="" type="radio"/> option 1 (preferred) <input type="radio"/> None		

If you cannot provide data on this indicator using the categories used in the proposed options, please send us a separate Excel table with data as reported in your country. In this Excel table please clearly specify the indicator, used categories and row and column headers.

### Core indicator #9: Distribution of parity

**Definition:** Number of previous live or stillbirths of women delivering a live or stillbirth.

*Attention: only include women delivering live and stillbirths at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☒ yes ☐ no

Can you provide data using another definition? ☐ yes ☐ no

Data source: 0

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of  
these options:

☒ option 1 (preferred)

☐ None

N women delivering  
live or stillborn babies

Parity

0

1

2

3

4+

Unknown

Calculated total

0



Core indicator #9: Distribution of parity		SAVE and return to INDEX
<b>Definition:</b> Number of previous live or stillbirths of women delivering a live or stillbirth. <b>Attention:</b> only include <u>women</u> delivering live and stillbirths at or after 22 completed weeks of gestation.		
Are you able to provide data using this definition? <input type="checkbox"/> yes <input type="checkbox"/> no Can you provide data using another definition? <input type="checkbox"/> yes <input type="checkbox"/> no		
<div></div>		To INDEX without saving
Data source: <input type="text" value="0"/> <input type="button" value="New data source"/>		
Please rate the quality of this indicator: <input checked="" type="radio"/> good <input type="radio"/> some concerns <input type="radio"/> bad		
Comments: <div></div>		Reset this page
Select one of these options:	<input checked="" type="radio"/> option 1 (preferred)	<input type="radio"/> None

If you cannot provide data on this indicator using the categories used in the proposed options, please send us a separate Excel table with data as reported in your country.  
 In this Excel table please clearly specify the indicator, used categories and row and column headers.

Core Indicator #10: Mode of delivery	
--------------------------------------	--

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☒ yes ☐ no

Are you able to provide data using this definition? ☒ yes ☐ no

Can you provide data using another definition? ☐ yes ☐ no

Can you provide data using another definition? ☐ yes ☐ no

Which definition is used for Caesarean sections: before labour/elective, during labour/emergency?

Data source: 0

New data source

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Comments:


SAVE and  
return to INDEX

To INDEX  
without saving

[Reset this page](#)

Select one of these options:

- option 1 (preferred)

 option 2

**option 3**

**option 4**

 option 5  c

Option 6 ☒ Non

ne

N live and stillbirths

Mode of delivery	Number of children	Percentage of children
Caesarean section	10	100%
Vaginal delivery	0	0%

Vaginal: non-instrumental

Vaginal: instrumental forceps

Vaginal: instrumental ventouse

Vaginal: instrumental other

Caesarean no labour/elective	
------------------------------	--

Caesarean in labour/emergency

Mode Unknown

Calculated total	0
------------------	---

Calculated total	0
------------------	---

Core indicator #10: Mode of delivery

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☐ yes
☐ no

Can you provide data using another definition?

☐ yes
☐ no

Which definition is used for Caesarean sections: before labour/elective, during labour/emergency?

SAVE and return to INDEX

To INDEX without saving

Data source: 0

New data source

Please rate the quality of this indicator:

☒ good
☐ some concerns
☐ bad

Comments:

Reset this page

Select one of these options:

☒ option 1 (preferred)
☐ option 2
☐ option 3
☐ option 4
☐ option 5
☐ option 6
☐ None

	N live and stillbirths
Mode of delivery	
Vaginal: non-instrumental	
Vaginal: instrumental	
Caesarean no labour/elective	
Caesarean in labour/emergency	
Mode Unknown	
Calculated total	0



Core indicator #10: Mode of delivery

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☐ yes
☐ no

Can you provide data using another definition?

☐ yes
☐ no

Which definition is used for Caesarean sections: before labour/elective, during labour/emergency?

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good
☐ some concerns
☐ bad

Comments:

SAVE and return to INDEX

To INDEX without saving

Reset this page

Select one of these options:

☒ option 1 (preferred)
☐ option 2
☐ option 3
☐ option 4
☐ option 5
☐ option 6
☐ None

	N live and stillbirths
Mode of delivery	
Vaginal	
Caesarean no labour/elective	
Caesarean in labour/emergency	
Mode Unknown	
Calculated total	0

Core Indicator #10: Mode of delivery

Definition: The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.

Are you able to provide data using this definition?

☐ yes
☐ no

Can you provide data using another definition?

☐ yes
☐ no

Which definition is used for Caesarean sections: before labour/elective, during labour/emergency?

Data source:

0

New data source

Please rate the quality of this indicator:

☐ good
☐ some concerns
☐ bad

Comments:

SAVE and return to INDEX

To INDEX without saving

Reset this page

Select one of these options:

☒ option 1 (preferred)
☐ option 2
☐ option 3
☐ option 4
☐ option 5
☐ option 6
☐ None

	N live and stillbirths
Mode of delivery	
Vaginal: non-instrumental	
Vaginal: instrumental forceps	
Vaginal: instrumental ventouse	
Vaginal: instrumental other	
Caesarean	
Mode Unknown	
Calculated total	0

Core Indicator #10: Mode of delivery

Definition: The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.

Are you able to provide data using this definition?

☐ yes
☐ no

Can you provide data using another definition?

☐ yes
☐ no

Which definition is used for Caesarean sections: before labour/elective, during labour/emergency?

Data source:

0

New data source

Please rate the quality of this indicator:

☐ good
☐ some concerns
☐ bad

Comments:

SAVE and return to INDEX

To INDEX without saving

Reset this page

Select one of these options:

☒ option 1 (preferred)
☐ option 2
☐ option 3
☐ option 4
☐ option 5
☐ option 6
☐ None

	N live and stillbirths
Mode of delivery	
Vaginal: non-instrumental	
Vaginal: instrumental	
Caesarean	
Mode Unknown	
Calculated total	0



Core indicator #10: Mode of delivery

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?
 

☒ yes
 ☐ no

Which definition is used for Caesarean sections: before labour/elective, during labour/emergency?
 

☐ yes
 ☐ no

Can you provide data using another definition?

Data source:
 

0

New data source

Please rate the quality of this indicator:
 

☒ good
 ☐ some concerns
 ☐ bad

Comments:

Select one of these options:
 

☒ option 1 (preferred)
 ☐ option 2
 ☐ option 3
 ☐ option 4
 ☐ option 5
 ☐ option 6
 ☐ None

N live and stillbirths

Mode of delivery
 

Vaginal
 
 Caesarean
 
 Mode Unknown
 
 Calculated total
 

0

SAVE and return to INDEX

To INDEX without saving

Reset this page

Core indicator #10: Mode of delivery		SAVE and return to INDEX
<b>Definition:</b> The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.		
<i>Attention: include all live and stillborn <u>babies</u> at or after 22 completed weeks of gestation.</i>		
Are you able to provide data using this definition? <input type="checkbox"/> yes <input type="checkbox"/> no Can you provide data using another definition? <input type="checkbox"/> yes <input type="checkbox"/> no	Which definition is used for Caesarean sections: before labour/elective, during labour/emergency?	To INDEX without saving
<div></div>		
Data source: <input type="text" value="0"/> <input type="button" value="New data source"/>		
Please rate the quality of this indicator: <input type="checkbox"/> good <input type="checkbox"/> some concerns <input type="checkbox"/> bad		
Comments: <input type="text"/>		Reset this page
Select one of these options: <input checked="" type="radio"/> option 1 (preferred) <input type="radio"/> option 2 <input type="radio"/> option 3 <input type="radio"/> option 4 <input type="radio"/> option 5 <input type="radio"/> option 6 <input type="radio"/> None		

If you cannot provide data on this indicator using the categories used in the proposed options, please send us a separate Excel table with data as reported in your country. In this Excel table please clearly specify the indicator, used categories and row and column headers.

Core indicator #10: Mode of delivery (by parity)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☒ yes
☐ no

Can you provide data using another definition?

☒ yes
☐ no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good
☐ some concerns
☐ bad

Comments:

SAVE and return to INDEX

To INDEX without saving

Reset this page

Select one of these options:

☒ option 1 (preferred)
☐ option 2
☐ option 3
☐ option 4
☐ option 5
☐ option 6
☐ None

Mode of delivery	----- Parity -----			Calc. totals
	Nullipara	Multipara	Parity Unknown	
Vaginal: non-instrumental				0
Vaginal: instrumental forceps				0
Vaginal: instrumental ventouse				0
Vaginal: instrumental other				0
Caesarean no labour/elective				0
Caesarean in labour/emergency				0
Mode Unknown				0
Calculated totals	0	0	0	0

Core indicator #10: Mode of delivery (by parity)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

**Attention:** include all live and stillborn babies at or after 22 completed weeks of gestation.

Are you able to provide data using this definition? ☐ yes ☐ no  
 Can you provide data using another definition? ☐ yes ☐ no

Data source: 0 New data source

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Comments:

SAVE and return to INDEX

To INDEX without saving

Reset this page

Select one of these options:

☒ option 1 (preferred)
 ☐ option 2
 ☐ option 3
 ☐ option 4
 ☐ option 5
 ☐ option 6
 ☐ None

	--- Parity ---			
	Nullipara	Multipara	Parity Unknown	Calc. totals
<b>Mode of delivery</b>				
Vaginal: non-instrumental				0
Vaginal: instrumental				0
Caesarean no labour/elective				0
Caesarean in labour/emergency				0
Mode Unknown				0
<b>Calculated totals</b>	0	0	0	0

**Core indicator #10: Mode of delivery (by parity)**

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☒ yes
☐ no

Can you provide data using another definition?

☐ yes
☒ no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good
☐ some concerns
☐ bad

Comments:

SAVE and return to INDEX

To INDEX without saving

Reset this page

Select one of these options:

☒ option 1 (preferred)
☐ option 2
☒ option 3
☐ option 4
☐ option 5
☐ option 6
☐ None

	Parity			
	Nullipara	Multipara	Parity Unknown	Calc. Totals
Mode of delivery				
Vaginal				0
Caesarean no labour/elective				0
Caesarean in labour/emergency				0
Mode Unknown				0
Calculated totals	0	0	0	0



Core indicator #10: Mode of delivery (by parity)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☐ yes
☐ no

Can you provide data using another definition?

☐ yes
☐ no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good
☐ some concerns
☐ bad

Comments:

SAVE and return to INDEX

To INDEX without saving

Reset this page

Select one of these options:

☒ option 1 (preferred)
☐ option 2
☐ option 3
☐ option 4
☐ option 5
☐ option 6
☐ None

	----- Nullipara	--- Parity --- Multipara	----- Parity Unknown	----- Calc. totals
<b>Mode of delivery</b>				
Vaginal: non-instrumental				0
Vaginal: instrumental forceps				0
Vaginal: instrumental ventouse				0
Vaginal: instrumental other				0
Caesarean				0
Mode Unknown				0
<i>Calculated totals</i>	0	0	0	0

### Core indicator #10: Mode of delivery (by parity)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☒ yes ☐ no

Can you provide data using another definition? ☐ yes ☒ no

Data source: 0 New data source

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☒ option 5

☐ option 6

☐ None

#### Mode of delivery

	Nullipara	Parity Multipara	Parity Unknown	Calc. totals
Vaginal: non-instrumental				0
Vaginal: instrumental				0
Caesarean				0
Mode Unknown				0
Calculated totals	0	0	0	0

### Core indicator #10: Mode of delivery (by parity)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☒ yes ☐ no

Can you provide data using another definition? ☐ yes ☐ no

Data source: 0 New data source

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

### Mode of delivery

	Nullipara	Parity Multipara	Parity Unknown	Calc. totals
Vaginal	0	0	0	0
Caesarean	0	0	0	0
Mode Unknown	0	0	0	0
Calculated totals	0	0	0	0

Core indicator #10: Mode of delivery (by parity)		SAVE and return to INDEX
<b>Definition:</b> The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.  <i>Attention: include all live and stillborn <u>babies</u> at or after 22 completed weeks of gestation.</i>		
Are you able to provide data using this definition? <input type="checkbox"/> yes <input type="checkbox"/> no Can you provide data using another definition? <input type="checkbox"/> yes <input type="checkbox"/> no		To INDEX without saving
<div> <div>Data source: 0</div> <div>New data source</div> </div> <div>           Please rate the quality of this indicator: <input type="checkbox"/> good <input type="checkbox"/> some concerns <input type="checkbox"/> bad         </div> <div>           Comments:         </div>		Reset this page
Select one of these options: <input checked="" type="radio"/> option 1 (preferred) <input type="radio"/> option 2 <input type="radio"/> option 3 <input type="radio"/> option 4 <input type="radio"/> option 5 <input type="radio"/> option 6 <input type="radio"/> None		

If you cannot provide data on this indicator using the categories used in the proposed options, please send us a separate Excel table with data as reported in your country.  
In this Excel table please clearly specify the indicator, used categories and row and column headers.



### Core indicator #10: Mode of delivery (by previous Caesarean Section)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☒ yes ☐ no

Can you provide data using another definition?

☒ yes ☐ no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good

☐ some concerns

☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

### Mode of delivery

	Previous CS			
	Yes	No	Unknown	Calc. Totals
Vaginal: non-instrumental				0
Vaginal: instrumental forceps				0
Vaginal: instrumental ventouse				0
Vaginal: instrumental other				0
Caesarean no labour/elective				0
Caesarean in labour/emergency				0
Mode Unknown				0
<b>Calculated totals</b>	0	0	0	0



### Core indicator #10: Mode of delivery (by previous Caesarean Section)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☒ yes

☐ no

Can you provide data using another definition?

☒ yes

☐ no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good

☐ some concerns

☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

### Mode of delivery

	Yes	No	Unknown	Calc. totals
Vaginal: non-instrumental				0
Vaginal: instrumental				0
Caesarean no labour/elective				0
Caesarean in labour/emergency				0
Mode Unknown				0
<b>Calculated totals</b>	0	0	0	0

Core indicator #10: Mode of delivery (by previous Caesarean Section)

Definition: The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.

Are you able to provide data using this definition?

☒ yes
☐ no

Can you provide data using another definition?

☐ yes
☐ no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good
☐ some concerns
☐ bad

Comments:

SAVE and return to INDEX

To INDEX without saving

Reset this page

Select one of these options:

☒ option 1 (preferred)
☐ option 2
☒ option 3
☐ option 4
☐ option 5
☐ option 6
☐ None

		Previous CS			
		Yes	No	Unknown	Calc. totals
Mode of delivery	Vaginal				0
	Caesarean no labour/elective				0
	Caesarean in labour/emergency				0
	Mode Unknown				0
	Calculated totals	0	0	0	0

**Core indicator #10: Mode of delivery (by previous Caesarean Section)**

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

**Attention:** include all live and stillborn babies at or after 22 completed weeks of gestation.

Are you able to provide data using this definition?

☒

no

Can you provide data using another definition?

☒

no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good

☐ some concerns

☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

**Mode of delivery**

	Yes	Previous CS No	Unknown	Calc. totals
Vaginal: non-instrumental				0
Vaginal: instrumental forceps				0
Vaginal: instrumental ventouse				0
Vaginal: instrumental other				0
Caesarean				0
Mode Unknown				0
<b>Calculated totals</b>	0	0	0	0

### Core indicator #10: Mode of delivery (by previous Caesarean Section)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☒

no

Can you provide data using another definition?

☒

no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good

☐ some concerns

☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

### Mode of delivery

Vaginal: non-instrumental

Vaginal: instrumental

Caesarean

Mode Unknown

Calculated totals

Yes

Previous CS  
No

Unknown

Calc. totals

0

0

0

0

0

0

0

0

0



### Core indicator #10: Mode of delivery (by previous Caesarean Section)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

**Attention:** include all live and stillborn babies at or after 22 completed weeks of gestation.

Are you able to provide data using this definition?

☐ yes

☐ no

Can you provide data using another definition?

☐ yes

☐ no

Data source:

0

New data source

Please rate the quality of this indicator:

☐ good

☐ some concerns

☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

### Mode of delivery

	Yes	Previous CS No	Unknown	Calc. totals
Vaginal				0
Caesarean				0
Mode Unknown				0
Calculated totals	0	0	0	0



Core indicator #10: Mode of delivery (by previous Caesarean Section)		SAVE and return to INDEX
<b>Definition:</b> The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.		
<i>Attention: include all live and stillborn <u>babies</u> at or after 22 completed weeks of gestation.</i>		
Are you able to provide data using this definition? <input type="checkbox"/> yes <input type="checkbox"/> no Can you provide data using another definition? <input type="checkbox"/> yes <input type="checkbox"/> no		To INDEX without saving
<div> <div></div> <div></div> </div>		
Data source: <input type="text" value="0"/> <span>New data source</span>		Reset this page
Please rate the quality of this indicator: <input type="checkbox"/> good <input type="checkbox"/> some concerns <input type="checkbox"/> bad		
Comments: <input type="text"/>		
Select one of these options: <input checked="" type="radio"/> option 1 (preferred) <input type="radio"/> option 2 <input type="radio"/> option 3 <input type="radio"/> option 4 <input type="radio"/> option 5 <input type="radio"/> option 6 <input type="radio"/> None		

If you cannot provide data on this indicator using the categories used in the proposed options, please send us a separate Excel table with data as reported in your country. In this Excel table please clearly specify the indicator, used categories and row and column headers.

### Core indicator #10: Mode of delivery (by presentation of fetus)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☒ yes ☐ no

Can you provide data using another definition? ☐ yes ☒ no

Data source:  New data source

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

### Mode of delivery

	Presentation				
	Breech	Vertex	- Other	-Unknown	Calc. totals
Vaginal: non-instrumental					0
Vaginal: instrumental forceps					0
Vaginal: instrumental ventouse					0
Vaginal: instrumental other					0
Caesarean no labour/elective					0
Caesarean in labour/emergency					0
Mode Unknown					0
Calculated totals	0	0	0	0	0

### Core indicator #10: Mode of delivery (by presentation of fetus)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☒

☐ no

Can you provide data using another definition?

☒

☐ no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good

☐ some concerns

☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

### Mode of delivery

	Breech	Vertex	- Other	-Unknown	Calc. totals
Vaginal: non-instrumental					0
Vaginal: instrumental					0
Caesarean no labour/elective					0
Caesarean in labour/emergency					0
Mode Unknown					0
<b>Calculated totals</b>	0	0	0	0	0

### Core indicator #10: Mode of delivery (by presentation of fetus)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☒ yes

☐ no

Can you provide data using another definition?

☐ yes

☐ no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good

☐ some concerns

☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

### Mode of delivery

	Presentation				
	Breech	Vertex	- Other	-Unknown	Calc. totals
Vaginal					0
Caesarean no labour/elective					0
Caesarean in labour/emergency					0
Mode Unknown					0
Calculated totals	0	0	0	0	0

### Core indicator #10: Mode of delivery (by presentation of fetus)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☒ yes

☐ no

Can you provide data using another definition?

☐ yes

☐ no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good

☐ some concerns

☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

### Mode of delivery

	Breech	Vertex	- Other	-Unknown	Calc. totals
Vaginal: non-instrumental					0
Vaginal: instrumental forceps					0
Vaginal: instrumental ventouse					0
Vaginal: instrumental other					0
Caesarean					0
Mode Unknown					0
Calculated totals	0	0	0	0	0



**Core indicator #10: Mode of delivery (by presentation of fetus)**

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☒

no

Can you provide data using another definition?

☒

no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good

☐ some concerns

☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☒ option 5

☐ option 6

☐ None

**Mode of delivery**

Vaginal: non-instrumental

Vaginal: instrumental

Caesarean

Mode Unknown

Calculated totals

Breech

Presentation  
Vertex

- Other

-Unknown

Calc. totals

0

0

0

0

0

0

0

0

0

### Core indicator #10: Mode of delivery (by presentation of fetus)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☒ yes

☐ no

Can you provide data using another definition?

☒ yes

☐ no

Data source:

0

New data source

Please rate the quality of this indicator:

☒ good

☐ some concerns

☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

Mode of delivery

	Breech	Presentation Vertex	- Other	-Unknown	Calc. totals
Vaginal					0
Caesarean					0
Mode Unknown					0
Calculated totals	0	0	0	0	0

Core Indicator #10: Mode of delivery (by presentation of fetus)		SAVE and return to INDEX
<b>Definition:</b> The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.  <i>Attention: include all live and stillborn <u>babies</u> at or after 22 completed weeks of gestation.</i>		
Are you able to provide data using this definition? <input checked="" type="checkbox"/> yes <input type="checkbox"/> no Can you provide data using another definition? <input type="checkbox"/> yes <input type="checkbox"/> no		To INDEX without saving
<div> <div>Data source: 0</div> <div>New data source</div> </div> <div>           Please rate the quality of this indicator: <input checked="" type="checkbox"/> good <input type="checkbox"/> some concerns <input type="checkbox"/> bad         </div> <div>           Comments:         </div>		Reset this page
Select one of these options: <input checked="" type="radio"/> option 1 (preferred) <input type="radio"/> option 2 <input type="radio"/> option 3 <input type="radio"/> option 4 <input type="radio"/> option 5 <input type="radio"/> option 6 <input type="radio"/> None		

If you cannot provide data on this indicator using the categories used in the proposed options, please send us a separate Excel table with data as reported in your country. In this Excel table please clearly specify the indicator, used categories and row and column headers.

Core indicator #10: Mode of delivery (by plurality)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☒ yes ☐ no  
Can you provide data using another definition? ☒ yes ☐ no

Data source: 0

New data source

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Comments:

SAVE and return to INDEX

To INDEX without saving

Reset this page

Select one of these options:

☒ option 1 (preferred)
☐ option 2
☐ option 3
☐ option 4
☐ option 5
☐ option 6
☐ None

	Plurality				
	Singleton	Twins	Triplets +	Plurality Unknown	Calc. totals
<b>Mode of delivery</b>					
Vaginal: non-instrumental					0
Vaginal: instrumental forceps					0
Vaginal: instrumental ventouse					0
Vaginal: instrumental other					0
Caesarean no labour/elective					0
Caesarean in labour/emergency					0
Mode Unknown					0
<i>Calculated totals</i>	0	0	0	0	0



### Core indicator #10: Mode of delivery (by plurality)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☒ yes ☐ no

Can you provide data using another definition? ☐ yes ☐ no

Data source: 0

New data source

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

### Mode of delivery

	Singleton	Plurality Twins	Triplets +	Plurality Unknown	Calc. totals
Vaginal: non-instrumental					0
Vaginal: instrumental					0
Caesarean no labour/elective					0
Caesarean in labour/emergency					0
Mode Unknown					0
<b>Calculated totals</b>	0	0	0	0	0



### Core Indicator #10: Mode of delivery (by plurality)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☒ yes ☐ no

Can you provide data using another definition? ☒ yes ☐ no

Data source: 0 New data source

Please rate the quality of this indicator: ☒ good ☐ some concerns ☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☒ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

Mode of delivery	Plurality				Calc. totals
	Singleton	Twins	Triplets +	Plurality Unknown	
Vaginal					0
Caesarean no labour/elective					0
Caesarean in labour/emergency					0
Mode Unknown					0
<b>Calculated totals</b>	0	0	0	0	0

### Core indicator #10: Mode of delivery (by plurality)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☐ yes ☐ no

Can you provide data using another definition? ☐ yes ☐ no

Data source: 0 New data source

Please rate the quality of this indicator: ☐ good ☐ some concerns ☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

#### Mode of delivery

	Singleton	Plurality Twins	Triplets +	Plurality Unknown	Calc. totals
Vaginal: non-instrumental					0
Vaginal: instrumental forceps					0
Vaginal: instrumental ventouse					0
Vaginal: instrumental other					0
Caesarean					0
Mode Unknown					0
<b>Calculated totals</b>	0	0	0	0	0

### Core indicator #10: Mode of delivery (by plurality)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition?

☐

no

Can you provide data using another definition?

☐

no

Data source:

0

New data source

Please rate the quality of this indicator:

☐

good

☐

some concerns

☐

bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒

option 1 (preferred)

☐

option 2

☐

option 3

☐

option 4

☒

option 5

☐

option 6

☐

None

### Mode of delivery

Vaginal: non-instrumental

Vaginal: instrumental

Caesarean

Mode Unknown

Calculated totals

Singleton

Plurality

Twins

Triplets +

Plurality Unknown

Calc. totals

0

0

0

0

0

0

0

0

0

0

**Core indicator #10: Mode of delivery** (by plurality)

**Definition:** The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.

*Attention: include all live and stillborn babies at or after 22 completed weeks of gestation.*

Are you able to provide data using this definition? ☐ yes ☐ no

Can you provide data using another definition? ☐ yes ☐ no

Data source: 0

Please rate the quality of this indicator: ☐ good ☐ some concerns ☐ bad

Comments:

SAVE and  
return to INDEX

To INDEX  
without saving

Reset this page

Select one of these  
options:

☒ option 1 (preferred)

☐ option 2

☐ option 3

☐ option 4

☐ option 5

☐ option 6

☐ None

**Mode of delivery**

	Singleton	Plurality Twins	Triplets +	Plurality Unknown	Calc. totals
Vaginal					0
Caesarean					0
Mode Unknown					0
<b>Calculated totals:</b>	0	0	0	0	0



Core indicator #10: Mode of delivery (by plurality)	
<p><b>Definition:</b> The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.</p> <p><i>Attention: include all live and stillborn <u>babies</u> at or after 22 completed weeks of gestation.</i></p>	SAVE and return to INDEX
<p>Are you able to provide data using this definition? <input type="checkbox"/> yes <input type="checkbox"/> no</p> <p>Can you provide data using another definition? <input type="checkbox"/> yes <input type="checkbox"/> no</p>	To INDEX without saving
<p>Data source: <input type="text" value="0"/> <input type="button" value="New data source"/></p> <p>Please rate the quality of this indicator: <input type="checkbox"/> good <input type="checkbox"/> some concerns <input type="checkbox"/> bad</p> <p>Comments: <input type="text"/></p>	Reset this page
<p>Select one of these options:</p> <div> <input checked="" type="radio"/> option 1 (preferred)           <input type="radio"/> option 2           <input type="radio"/> option 3           <input type="radio"/> option 4           <input type="radio"/> option 5           <input type="radio"/> option 6           <input type="radio"/> None         </div>	

If you cannot provide data on this indicator using the categories used in the proposed options, please send us a separate Excel table with data as reported in your country. In this Excel table please clearly specify the indicator, used categories and row and column headers.



















Core indicator #10: Mode of delivery (by gestational age)		SAVE and return to INDEX
<b>Definition:</b> The number of live and stillbirths associated with each of the following modes of delivery: Vaginal (non-instrumental, instrumental forceps/ventouse/other, such as Thierry's spatula), Caesarean undertaken before or at onset of labour/elective, Caesarean undertaken during labour/emergency, as a proportion of all live and stillbirths.  <i>Attention: include all live and stillborn <u>babies</u> at or after 22 completed weeks of gestation.</i>		
Are you able to provide data using this definition? <input type="checkbox"/> yes <input type="checkbox"/> no Can you provide data using another definition? <input type="checkbox"/> yes <input type="checkbox"/> no		To INDEX without saving
<div> <div>Data source: 0</div> <div>New data source</div> </div> <div>           Please rate the quality of this indicator: <input type="checkbox"/> good <input type="checkbox"/> some concerns <input type="checkbox"/> bad         </div> <div>           Comments:         </div>		
Select one of these options: <input type="checkbox"/> option 1 (preferred) <input type="checkbox"/> option 2 <input type="checkbox"/> option 3 <input type="checkbox"/> option 4 <input type="checkbox"/> option 5 <input type="checkbox"/> option 6 <input type="checkbox"/> None		Reset this page

If you cannot provide data on this indicator using the categories used in the proposed options, please send us a separate Excel table with data as reported in your country. In this Excel table please clearly specify the indicator, used categories and row and column headers.